







- [8] SATI S, GHOSH S, JAIN V, SCARIA V, SENGUPTA S. Genome-wide analysis reveals distinct patterns of epigenetic features in long non-coding RNA loci[J]. *Nucleic Acids Res*, 2012, 40: 10018-10031.
- [9] MARTIANOV I, RAMADASS A, SERRA BARROS A, CHOW N, AKOULITCHEV A. Repression of the human dihydrofolate reductase gene by a non-coding interfering transcript[J]. *Nature*, 2007, 445: 666-670.
- [10] RINN J L, KERTESZ M, WANG J K, SQUAZZO S L, XU X, BRUGMANN S A, et al. Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs[J]. *Cell*, 2007, 129: 1311-1323.
- [11] SHAH N, SUKUMAR S. The *Hox* genes and their roles in oncogenesis[J]. *Nat Rev Cancer*, 2010, 10: 361-371.
- [12] ENDO H, SHIROKI T, NAKAGAWA T, YOKOYAMA M, TAMAI K, YAMANAMI H, et al. Enhanced expression of long non-coding RNA HOTAIR is associated with the development of gastric cancer[J/OL]. *PLoS One*, 2013. doi:10.1371/journal.pone.0077071
- [13] XU Z Y, YU Q M, DU Y A, YANG L T, DONG R Z, HUANG L, et al. Knockdown of long non-coding RNA HOTAIR suppresses tumor invasion and reverses epithelial-mesenchymal transition in gastric cancer[J]. *Int J Biol Sci*, 2013, 9: 587-597.
- [14] LIU Y W, SUN M, XIA R, ZHANG E B, LIU X H, ZHANG Z H, et al. LincHOTAIR epigenetically silences miR34a by binding to PRC2 to promote the epithelial-to-mesenchymal transition in human gastric cancer[J/OL]. *Cell Death Dis*, 2015, 6: e1802. doi:10.1038/cddis.2015.150.
- [15] LOTTIN S, ADRIAENSSENS E, DUPRESSOIR T, BERTEAUX N, MONTPELLIER C, COLL J, et al. Overexpression of an ectopic *H19* gene enhances the tumorigenic properties of breast cancer cells[J]. *Carcinogenesis*, 2002, 23: 1885-1895.
- [16] ZHANG L, YANG F, YUAN J H, YUAN S X, ZHOU W P, HUO X S, et al. Epigenetic activation of the miR-200 family contributes to H19-mediated metastasis suppression in hepatocellular carcinoma[J]. *Carcinogenesis*, 2013, 34: 577-586.
- [17] BERTEAUX N, LOTTIN S, MONTÉ D, PINTE S, QUATANENS B, COLL J, et al. H19 mRNA-like noncoding RNA promotes breast cancer cell proliferation through positive control by E2F1[J]. *J Biol Chem*, 2005, 280: 29625-29636.
- [18] MATOUK I J, RAVEH E, ABU-LAIL R, MEZAN S, GILON M, GERSHTAIN E, et al. Oncofetal H19 RNA promotes tumor metastasis[J]. *Biochim Biophys Acta*, 2014, 1843: 1414-1426.
- [19] ZHOU X, YE F, YIN C, ZHUANG Y, YUE G, ZHANG G. The interaction between miR-141 and lncRNA-H19 in regulating cell proliferation and migration in gastric cancer[J]. *Cell Physiol Biochem*, 2015, 36: 1440-1452.
- [20] SAITO T, KURASHIGE J, NAMBARA S, KOMATSU H, HIRATA H, UEDA M, et al. A long non-coding RNA activated by transforming growth factor-beta is an independent prognostic marker of gastric cancer[J]. *Ann Surg Oncol*, 2015, 22: 915-922.
- [21] LEI K, LIANG X, GAO Y, XU B, XU Y, LI Y, et al. Lnc-ATB contributes to gastric cancer growth through a miR-141-3p/TGFβ2 feedback loop[J]. *Biochem Biophys Res Commun*, 2017, 484: 514-521.
- [22] LING H, SPIZZO R, ATLASI Y, NICOLOSO M, SHIMIZU M, REDIS R S, et al. CCAT2, a novel noncoding RNA mapping to 8q24, underlies metastatic progression and chromosomal instability in colon cancer[J]. *Genome Res*, 2013, 23: 1446-1461.
- [23] WANG Y J, LIU J Z, LV P, DANG Y, GAO J Y, WANG Y. Long non-coding RNA CCAT2 promotes gastric cancer proliferation and invasion by regulating the E-cadherin and LATS2[J]. *Am J Cancer Res*, 2016, 6: 2651-2660.
- [24] YU Y, LI L, ZHENG Z, CHEN S, CHEN E, HU Y. Long non-coding RNA linc00261 suppresses gastric cancer progression via promoting Slug degradation[J]. *J Cell Mol Med*, 2017, 21: 955-967.
- [25] HAN Y, YE J, WU D, WU P, CHEN Z, CHEN J, et al. LEIGC long non-coding RNA acts as a tumor suppressor in gastric carcinoma by inhibiting the epithelial-to-mesenchymal transition[J/OL]. *BMC Cancer*, 2014, 14: 932. doi: 10.1186/1471-2407-14-932.
- [26] ZHAO J, LIU Y, ZHANG W, ZHOU Z, WU J, CUI P, et al. Long non-coding RNA Linc00152 is involved in cell cycle arrest, apoptosis, epithelial to mesenchymal transition, cell migration and invasion in gastric cancer[J]. *Cell Cycle*, 2015, 14: 3112-3123.
- [27] XIE M, NIE F Q, SUN M, XIA R, LIU Y W, ZHOU P, et al. Decreased long noncoding RNA SPRY4-IT1 contributing to gastric cancer cell metastasis partly via affecting epithelial-mesenchymal transition[J/OL]. *J Transl Med*, 2015, 13: 250. doi: 10.1186/s12967-015-0595-9.
- [28] ZHOU H, WANG F, CHEN H, TAN Q, QIU S, CHEN S, et al. Increased expression of long-noncoding RNA ZFAS1 is associated with epithelial-mesenchymal transition of gastric cancer[J]. *Aging (Albany NY)*, 2016, 8: 2023-2035.
- [29] ARUMUGAM T, RAMACHANDRAN V, FOURNIER K F, WANG H, MARQUIS L, ABBRUZZESE J L, et al. Epithelial to mesenchymal transition contributes to drug resistance in pancreatic cancer[J]. *Cancer Res*, 2009, 69: 5820-5828.
- [30] BLICK T, HUGO H, WIDODO E, WALTHAM M, PINTO C, MANI S A, et al. Epithelial mesenchymal transition traits in human breast cancer cell lines parallel the CD44<sup>hi</sup>CD24<sup>lo/-</sup> stem cell phenotype in human breast cancer[J]. *J Mammary Gland Biol Neoplasia*, 2010, 15: 235-252.
- [31] FISCHER K R, DURRANS A, LEE S, SHENG J, LI F, WONG S T, et al. Epithelial-to-mesenchymal transition is not required for lung metastasis but contributes to chemoresistance[J]. *Nature*, 2015, 527: 472-476.